



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MECHANICS.

344. Proposed by J. ROSENBAUM, New Haven, Conn.

Two bodies of equal masses, and coefficients of friction μ_1 and μ_2 are connected by a light, flexible string, and placed on an inclined plane. What is the angle, θ , between the string and the plane if the inclination, α , of the plane is a minimum when the bodies are on the point of motion?

345. Proposed by J. L. RILEY, Northeastern State Normal School, Tahlequah, Okla.

Two particles A and B are together in a smooth circular tube. A attracts B with a force whose acceleration is ω^2 and moves along the tube with uniform angular velocity 2ω , B being initially at rest: prove that the angle Φ subtended by AB at the center after a time t is given by the equation

$$\log \tan \frac{\pi + \Phi}{4} = \omega t.$$

NUMBER THEORY.

263. Proposed by J. L. RILEY, Northeastern State Normal School, Tahlequah, Okla.

To find values, positive integral, which verify the equation.

$$X^3 + 2 = Y^2 \quad (\text{Gerono}).$$

264. Proposed by C. F. GUMMER, Kingston, Ontario.

Find a general formula for three squares in arithmetical progression. Is it possible for the common difference to be a perfect square?

SOLUTIONS OF PROBLEMS.

ALGEBRA.

A solution of 464, by Mrs. ELIZABETH BROWN DAVIS, was received after selections had been made for publication.

455. Proposed by JOS. B. REYNOLDS, Lehigh University.

Solve for x_n (not in determinant form) the simultaneous equations,

$$\frac{4}{3}x_n + 2x_{n-1} + 2x_{n-2} \cdots 2x_4 + 2x_3 + 2x_2 + 2x_1 = g,$$

$$\frac{10}{3}x_n + \frac{22}{3}x_{n-1} + 8x_{n-2} \cdots 8x_4 + 8x_3 + 8x_2 + 8x_1 = 4g,$$

$$\frac{1}{3}x_n + \frac{4}{3}x_{n-1} + \frac{5}{3}x_{n-2} \cdots 18x_4 + 18x_3 + 18x_2 + 18x_1 = 9g,$$

$$\frac{22}{8}x_n + \frac{58}{8}x_{n-1} + \frac{82}{8}x_{n-2} + \frac{94}{8}x_{n-3} \cdots 32x_4 + 32x_3 + 32x_2 + 32x_1 = 16g,$$

$$\frac{28}{3}x_n + \frac{76}{3}x_{n-1} + \frac{112}{3}x_{n-2} + \frac{136}{3}x_{n-3} + \frac{148}{3}x_{n-4} \cdots = 25g,$$

$$(\tfrac{1}{3} + 2n - 1)x_n + (\tfrac{1}{3} + 6n - 5)x_{n-1} + (\tfrac{1}{3} + 10n - 13)x_{n-2} + (\tfrac{1}{3} + 14n - 25)x_{n-3}$$

$$\cdots (\tfrac{1}{3} + 2n^2 - 1)x_1 = n^2 g.$$

SOLUTION BY C. F. GUMMER, Kingston, Ont.

The equations may be written

$$2x_n + 3x_{n-1} + 3x_{n-2} + \cdots + 3x_1 = \frac{3g}{2},$$

$$5x_n + 11x_{n-1} + 12x_{n-2} + \cdots + 12x_1 = \frac{12g}{2},$$

$$8x_n + 20x_{n-1} + 26x_{n-2} + 27x_{n-3} + \cdots + 27x_1 = \frac{27g}{2},$$

$$\begin{array}{cccccccccccccccc} \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ (3n-1)x_n + (9n-7)x_{n-1} + \cdots + (3n^2-1)x_1 = \frac{3n^2g}{2}, \end{array}$$